

機密 (只限閱卷員使用)
CONFIDENTIAL (FOR MARKER'S USE ONLY)

香港考試及評核局
HONG KONG EXAMINATIONS AND ASSESSMENT AUTHORITY

2022年香港中學文憑考試
HONG KONG DIPLOMA OF SECONDARY EDUCATION EXAMINATION 2022

數學 **必修部分** **試卷一**
MATHEMATICS **COMPULSORY PART** **PAPER 1**

評卷參考
MARKING SCHEME

本評卷參考乃香港考試及評核局專為今年本科考試而編寫，供閱卷員參考之用。本評卷參考之使用，均受制於閱卷員有關之服務合約條款及閱卷員指引。特別是：

- 本局擁有並保留本評卷參考的所有財產權利(包括知識產權)。在未獲本局之書面批准下，閱卷員均不得複製、發表、透露、提供、使用或經營本評卷參考之全部或其部份。在遵守上述條款之情況下，本局有限地容許閱卷員可在應屆香港中學文憑考試的考試成績公布後，將本評卷參考提供任教本科的教師參閱。
- 在任何情況下，均不得容許本評卷參考之全部或其部份落入學生手中。本局籲請各閱卷員/教師通力合作，堅守上述原則。

This marking scheme has been prepared by the Hong Kong Examinations and Assessment Authority for the reference of markers. The use of this marking scheme is subject to the relevant service agreement terms and Instructions to Markers. In particular:

- The Authority retains all proprietary rights (including intellectual property rights) in this marking scheme. This marking scheme, whether in whole or in part, must not be copied, published, disclosed, made available, used or dealt in without the prior written approval of the Authority. Subject to compliance with the foregoing, a limited permission is granted to markers to share this marking scheme, after release of examination results of the current HKDSE examination, with teachers who are teaching the same subject.
- Under no circumstances should students be given access to this marking scheme or any part of it. The Authority is counting on the co-operation of markers/teachers in this regard.



機密 (只限閱卷員使用)
CONFIDENTIAL (FOR MARKER'S USE ONLY)

Hong Kong Diploma of Secondary Education Examination
Mathematics Compulsory Part Paper 1

General Marking Instructions

1. It is very important that all markers should adhere as closely as possible to the marking scheme. In many cases, however, candidates will have obtained a correct answer by an alternative method not specified in the marking scheme. In general, a correct answer merits *all the marks* allocated to that part, unless a particular method has been specified in the question. Markers should be patient in marking alternative solutions not specified in the marking scheme.
2. In the marking scheme, marks are classified into the following three categories:

‘M’ marks	awarded for correct methods being used;
‘A’ marks	awarded for the accuracy of the answers;
Marks without ‘M’ or ‘A’	awarded for correctly completing a proof or arriving at an answer given in a question.

In a question consisting of several parts each depending on the previous parts, ‘M’ marks should be awarded to steps or methods correctly deduced from previous answers, even if these answers are erroneous. However, ‘A’ marks for the corresponding answers should NOT be awarded (unless otherwise specified).
3. For the convenience of markers, the marking scheme was written as detailed as possible. However, it is still likely that candidates would not present their solution in the same explicit manner, e.g. some steps would either be omitted or stated implicitly. In such cases, markers should exercise their discretion in marking candidates’ work. In general, marks for a certain step should be awarded if candidates’ solution indicated that the relevant concept/technique had been used.
4. In marking candidates’ work, the benefit of doubt should be given in the candidates’ favour.
5. In the marking scheme, ‘r.t.’ stands for ‘accepting answers which can be rounded off to’ and ‘f.t.’ stands for ‘follow through’. Steps which can be skipped are shaded whereas alternative answers are enclosed with rectangles. All fractional answers must be simplified.

Solution	Marks	Remarks
1. $\frac{(a^3b^{-2})^4}{a^{-5}b^6}$ $= \frac{a^{12}b^{-8}}{a^{-5}b^6}$ $= \frac{a^{12+5}}{b^{6+8}}$ $= \frac{a^{17}}{b^{14}}$	1M 1M 1A -----(3)	for $(m^h)^k = m^{hk}$ or $(mn)^l = m^l n^l$ for $\frac{x^p}{x^q} = x^{p-q}$ or $y^{-r} = \frac{1}{y^r}$
2. Note that $x + y = 456$ and $7x = y$. So, we have $x + 7x = 456$. Solving, we have $x = 57$.	1A 1M 1A	for either correct for getting a linear equation in x or y only
$\frac{x}{1+7}$ $= \frac{456}{8}$ $= 57$	1M+1A 1A -----(3)	1M for fraction
3. $\frac{3}{k-9} + \frac{2}{5k+6}$ $= \frac{3(5k+6) + 2(k-9)}{(k-9)(5k+6)}$ $= \frac{15k+18+2k-18}{(k-9)(5k+6)}$ $= \frac{17k}{(k-9)(5k+6)}$	1M 1M 1A -----(3)	or equivalent
4. (a) $9c^2 - 6c + 1$ $= (3c-1)^2$ (b) $(4c+d)^2 - 9c^2 + 6c - 1$ $= (4c+d)^2 - (9c^2 - 6c + 1)$ $= (4c+d)^2 - (3c-1)^2$ $= (4c+d+3c-1)(4c+d-(3c-1))$ $= (7c+d-1)(c+d+1)$	1A 1M 1M 1A -----(4)	or equivalent for using the result of (a) or equivalent

Solution	Marks	Remarks
<p>5. Let \$ x \$ be the cost of the fan.</p> $(26\%)x = 78$ $x = \frac{78}{0.26}$ $x = 300$ <p>The selling price of the fan</p> $= 300 + 78$ $= \$ 378$ <p>Let \$ y \$ be the marked price of the fan.</p> $(70\%)y = 378$ $y = \frac{378}{0.7}$ $y = 540$ <p>Thus, the marked price of the fan is \$ 540 .</p>	<p>1M</p> <p>1M</p> <p>1M</p> <p>1A</p>	
<div style="border: 1px solid black; padding: 5px;"> <p>The marked price of the fan</p> $= \frac{(78)(1+26\%)}{(26\%)(70\%)}$ $= \\$ 540$ </div>	<p>1M+1M+1M</p> <p>1A</p>	<p>{ 1M for fraction + 1M for numerator + 1M for denominator</p>
------(4)		
<p>6. (a) $-2(3x+2) > x+10$</p> $-6x-4 > x+10$ $-6x-x > 10+4$ $-7x > 14$ $x < -2$ $2x \leq -8$ $x \leq -4$ <p>Therefore, we have $x < -2$ or $x \leq -4$.</p> <p>Thus, the solution of (*) is $x < -2$.</p>	<p>1M</p> <p>1A</p> <p>1A</p>	<p>for putting x on one side</p>
<p>(b) -3</p>	<p>1A</p>	
------(4)		
<p>7. (a) The coordinates of S' are $(5, 12)$.</p> <p>The coordinates of T' are $(-3, 7)$.</p>	<p>1A</p> <p>1A</p>	
<p>(b) The slope of $S'T'$</p> $= \frac{12-7}{5-(-3)}$ $= \frac{5}{8}$	<p>1M</p> <p>1A</p>	<p>0.625</p>
------(4)		

	Solution	Marks	Remarks
8. (a)	$\angle ACB = \angle CAD$ (alt. \angle s, $AD \parallel BC$) $\angle CAD = \angle ADE$ (alt. \angle s, $AC \parallel ED$) $\angle ACB = \angle ADE$ $\angle ABC = \angle AED$ (given) $AB = AE$ (given) $\triangle ABC \cong \triangle AED$ (AAS)		[(內)錯角, $AD \parallel BC$] [(內)錯角, $AC \parallel ED$] [已知] [已知]
Marking Scheme:			
	Case 1 Any correct proof with correct reasons.	2	
	Case 2 Any correct proof without reasons.	1	
(b)	$\angle BAC$ $= \angle DAE$ $= 87^\circ$ $\angle ACB$ $= 180^\circ - \angle BAC - \angle ABC$ $= 180^\circ - 87^\circ - 39^\circ$ $= 54^\circ$ $\angle CAD$ $= \angle ACB$ $= 54^\circ$ Note that $AC = AD$. $\angle ACD$ $= \angle ADC$ $= \frac{180^\circ - \angle CAD}{2}$ $= \frac{180^\circ - 54^\circ}{2}$ $= 63^\circ$	1M 1M 1A -----(5)	
9. (a)	12	1A	
(b)	Note that $a = 3$ and $b = 5$. The mean $= \frac{12(3) + 17(9) + 22(5) + 27(3)}{20}$ $= 19$ minutes	1M 1A	
(c)	The required probability $= \frac{12}{20}$ $= \frac{3}{5}$	1M 1A -----(5)	for numerator 0.6

Solution	Marks	Remarks
10. (a) Let $f(x) = ax^2 + bx$, where a and b are non-zero constants. So, we have $16a + 4b = 96$ and $25a - 5b = 15$. Solving, we have $a = 3$ and $b = 12$. Thus, we have $f(x) = 3x^2 + 12x$.	1A 1M 1A	for either substitution for both correct
----- (3)		
(b) The x -intercepts of the graph of $y = 8f(x)$ are 0 and -4 .	1A	for both correct
----- (1)		
(c) The equation $3x^2 + 12x - k = 0$ has two distinct real roots. $12^2 - 4(3)(-k) > 0$ $k > -12$	1M 1A	
----- (2)		
11. (a) $36 - (20 + a) = 14$ $a = 2$ $30 + b = 31$ $b = 1$	1M 1A 1A	
----- (3)		
(b) (i) The original mode $= 36$ The new mode $= 36$ Thus, there is no change in the mode of the distribution.	1M 1A	----- either one ----- f.t.
(ii) There are two cases. Case 1: The player of age 17 leaves the football team. The standard deviation of the distribution ≈ 7.162537194 Case 2: The player of age 43 leaves the football team. The standard deviation of the distribution ≈ 7.132307207 Thus, the greatest possible standard deviation of the distribution is 7.16.	1M 1A	----- either one ----- f.t.
----- (4)		

Solution	Marks	Remarks
13. (a) The ratio of the volume of the smaller sphere to the volume of the larger sphere is 8 : 27 .	1M	
The volume of the smaller sphere		
$= \frac{4}{3} \pi (9)^3 \left(\frac{8}{27} \right)$	1M	
$= 288\pi \text{ cm}^3$	1A	
	----- (3)	
(b) The volume of A		
$= \frac{1}{3} \pi (6^2)(10)$	1M	
$= 120\pi \text{ cm}^3$		
The volume of B		
$= 288\pi + \frac{4}{3} \pi (9)^3 - 120\pi$	1M	
$= 1140\pi \text{ cm}^3$		
$\frac{\text{The volume of } B}{\text{The volume of } A}$		
$= \frac{1140\pi}{120\pi}$	1M	
$= \frac{19}{2}$		
$\left(\frac{\text{The base radius of } B}{\text{The base radius of } A} \right)^3$		either one
$= \left(\frac{12}{6} \right)^3$		
$= 8$		
$\neq \frac{19}{2}$		
So, A and B are not similar.		
Thus, the claim is not correct.	1A	f.t.
	----- (4)	

Solution	Marks	Remarks
14. (a) Let $p(x) = (mx + n)(x^2 - 2x + 3) + x + 13$, where m and n are constants. Therefore, we have $p(x) = mx^3 + (n - 2m)x^2 + (3m - 2n + 1)x + 3n + 13$. So, we have $m = 2$ and $3n + 13 = -20$. Solving, we have $m = 2$ and $n = -11$. Hence, we have $p(x) = 2x^3 - 15x^2 + 29x - 20$. Thus, we have $a = -15$ and $b = 29$.	1M 1M 1A	for either one for both correct
(b) $p(5)$ $= 2(5)^3 - 15(5)^2 + 29(5) - 20$ $= 0$ Thus, $x - 5$ is a factor of $p(x)$.	1M 1A	f.t.
(c) $p(x) = 0$ $(x - 5)(2x^2 - 5x + 4) = 0$ $x - 5 = 0$ or $2x^2 - 5x + 4 = 0$ $(-5)^2 - 4(2)(4)$ $= -7$ < 0 So, the quadratic equation $2x^2 - 5x + 4 = 0$ has no real roots. Hence, the quadratic equation $2x^2 - 5x + 4 = 0$ has no irrational roots. Note that 5 is not an irrational number. Therefore, the equation $p(x) = 0$ has no irrational roots. Thus, the claim is disagreed.	1M 1M 1A	f.t.

Solution	Marks	Remarks
15. (a) The required probability $= \frac{C_2^{10} C_2^{12}}{C_4^{22}}$ $= \frac{54}{133}$	1M 1A	for numerator r.t. 0.406
The required probability $= 6 \left(\frac{10}{22} \right) \left(\frac{9}{21} \right) \left(\frac{12}{20} \right) \left(\frac{11}{19} \right)$ $= \frac{54}{133}$	1M 1A	for $6p_1 p_2 p_3 p_4$ r.t. 0.406
(b) The required probability $= 1 - \frac{54}{133}$ $= \frac{79}{133}$	-----(2) 1M 1A	for $1 - (a)$ r.t. 0.594
The required probability $= \frac{C_4^{10}}{C_4^{22}} + \frac{C_3^{10} C_1^{12}}{C_4^{22}} + \frac{C_1^{10} C_3^{12}}{C_4^{22}} + \frac{C_4^{12}}{C_4^{22}}$ $= \frac{79}{133}$	1M 1A	for $p_5 + p_6 + p_7 + p_8$ r.t. 0.594
16. (a) $g(x)$ $= 3x^2 + 12kx + 16k^2 + 8$ $= 3(x^2 + 4kx) + 16k^2 + 8$ $= 3(x^2 + 4kx + 4k^2) + 4k^2 + 8$ $= 3(x + 2k)^2 + 4k^2 + 8$ Thus, the coordinates of the vertex are $(-2k, 4k^2 + 8)$.	1M 1A ----- (2)	
(b) The coordinates of B are $(2k, 8k^2 + 16)$. Note that $AM : MB = 1 : 3$.	1M	
The coordinates of M $= \left(\frac{3(-2k) + (2k)}{1+3}, \frac{3(4k^2 + 8) + (8k^2 + 16)}{1+3} \right)$ $= (-k, 5k^2 + 10)$	1M 1A	----- (3)

機密 (只限閱卷員使用)
CONFIDENTIAL (FOR MARKER'S USE ONLY)

Solution	Marks	Remarks
17. (a) Note that $\alpha + \beta = -c$ and $\alpha\beta = -9$. $\alpha^2 + \beta^2$ $= (\alpha + \beta)^2 - 2\alpha\beta$ $= (-c)^2 - 2(-9)$ $= c^2 + 18$	1M 1M 1A	for either one
$c^2 + 18 - c^2 = 85 - (c^2 + 18)$ $c^2 = 49$ <p>Note that the 1st term and the common difference of the sequence are 49 and 18 respectively.</p> $\frac{n}{2}(2(49) + 18(n-1)) > 2 \times 10^6$ $9n^2 + 40n - 2 \times 10^6 > 0$ $n < \frac{-40 - \sqrt{40^2 - 4(9)(-2 \times 10^6)}}{2(9)} \quad \text{or} \quad n > \frac{-40 + \sqrt{40^2 - 4(9)(-2 \times 10^6)}}{2(9)}$ $n < -473.6319808 \quad \text{or} \quad n > 469.1875364$ Thus, the least value of n is 470 .	----- (3) 1M 1M 1A	
	----- (4)	

Solution	Marks	Remarks
18. (a) (i) $QR^2 = PQ^2 + PR^2 - 2(PQ)(PR)\cos\angle QPR$ $QR^2 = 30^2 + 25^2 - 2(30)(25)\cos 95^\circ$ $QR \approx 40.69070673$ $QR \approx 40.7$ cm Thus, the length of QR is 40.7 cm .	1M 1A	 r.t. 40.7 cm
(ii) $\frac{\sin\angle PQR}{PR} = \frac{\sin\angle QPR}{QR}$ $\frac{\sin\angle PQR}{25} \approx \frac{\sin 95^\circ}{40.69070673}$ $\angle PQR \approx 37.73809375^\circ$ or $\angle PQR \approx 142.2619063^\circ$ (rejected) Thus, we have $\angle PQR \approx 37.7^\circ$.	1M 1A	 r.t. 37.7°
----- (4)		
(b) $PM^2 = PQ^2 + QM^2 - 2(PQ)(QM)\cos\angle PQR$ $PM^2 \approx 30^2 + \left(\frac{40.69070673}{2}\right)^2 - 2(30)\left(\frac{40.69070673}{2}\right)\cos 37.73809375^\circ$ $PM \approx 18.66993831$ cm Let D and N be the projections of R and M on the horizontal ground respectively.		
MN $= \frac{1}{2}RD$ $= \frac{1}{2}PR\sin 70^\circ$ $= \frac{1}{2}(25)\sin 70^\circ$ ≈ 11.74615776 cm	1M	
Note that the angle between PM and the horizontal ground is $\angle MPN$. $\sin\angle MPN = \frac{MN}{PM}$ $\sin\angle MPN \approx \frac{11.74615776}{18.66993831}$ $\angle MPN \approx 38.98730493^\circ$ $\angle MPN < 40^\circ$ Thus, the claim is not correct.	1M 1A	 f.t.
----- (3)		

Solution	Marks	Remarks
<p>(d) Note that $\angle APG = \angle AQG = 90^\circ$ and $\angle APG + \angle AQG = 180^\circ$. So, $APGQ$ is a cyclic quadrilateral and AG is a diameter of the circumcircle of $\triangle APQ$. The radius of the circumcircle of $\triangle APQ$ $= \frac{1}{2} \sqrt{(83 - 158)^2 + (112 - 12)^2}$ $= \frac{125}{2}$</p> <p>By (c), the radius of the inscribed circle of $\triangle APQ$ is 30 .</p> <p>The ratio of the area of the inscribed circle to the area of the circumcircle of $\triangle APQ$ $= 30^2 : \left(\frac{125}{2}\right)^2$ $= 144 : 625$ $\neq 1 : 4$ Thus, the claim is disagreed.</p>	<p>1M</p> <p>1M</p> <p>1A ----- (3)</p>	<p>f.t.</p>

Paper 2

Since 1996

MC answer									
1	2	3	4	5	6	7	8	9	10
D	B	A	D	A	B	A	B	D	A
C	C	B	C	A	D	B	C	D	B
B	C	C	A	D	A	C	D	A	B
C	C	D	A	B	B	D	D	C	A
A	B	B	C	D					